

УДК 616.248-06

**THE OCCUR DEGREE OF THE BRONCHOOBSTRUCTIVE SYNDROME'S  
DEPENDENCE TO THE HUE OF THE REFLUXATE IN THE  
GASTROESOPHAGEAL REFLUX DISEASE****Yuldasheva D.H.***Bukhara State Medical Institute. City Bukhara. Uzbekistan*

The urgency of the problem. Nowadays, gastroesophageal reflux disease (GERD) holds the high positions according to happening index among the gastroenterological diseases. The diagnosing disease, almost is not difficult, because typical clinical symptoms of the GERD observed frequently in the clinical practice, for example heartburn, burp, regurgitating and others. However, typical forms of the disease are diagnosed much difficulty. Because clinical view of them looks like the diseases of the neighbor organs [1,2,4,6,7]. The broncho-pulmonary form of the GERD from atypical views is much observed. For instance, according to medical literatures one of atypical forms of the GERD reflux –asthma from 30 % until 90 % happens together with bronchial asthma (BA) [2,3,5,8,9].

According to our primary researchings among the atypical forms of the GERD, precisely, broncho-pulmonary type is progressive because we aimed during the research to define the immixture of register grade of the bronchoobstructive syndrome (BOS) with hue of the refluxate.

Materials and methods. The clinical, laboratory and instrumental facts of analysis results of 86 seconded patients, which are diseased with GERD and alternatively diseased with 1-2 steps of BOS were taken for the research. From the patients 51 women and 35 men ages between 18 and 57 (medium 35 + 0,7). Estimating benchmarks to the exploration: they are those people who have complaints about heartburn and regurgitation, older from 18 age and agreed in writing form for the clinical and instrumental analyzing. Releasing benchmarks from the exploration: consumption H2 blockers, proton pump inhibitors, prokinetics before 10 days addressing, attack level of the ulcer stomach and duodenum, achalazia of cardia, cirrhosis of the liver in various etiology, chronic pancreatitis in the twinge level, cholecystitis, choledocholithiasis disease, infection *Helicobacter pylori*, in the anamnesis operations were thereupon bile, pancreatic and duodenum diseases, pregnant and in the lactation period in the anamnesis which drugs were necessary underlined adverse effects or useless drugs and injuring with chronic deficit of ren.

For the diagnosing used from rentgenological, endoscopic and in necessary times ultrasonography and computer tomography examinations. Types of the refluxate ambience were determined, which were invited by authors in vivo and in vitro haller in the early morning first quality, then quantity with ph-metric method.

Before doing this method, patients were invited to reject juices, coffee and antacid drugs. We used classification that was offered in 2009 during the diagnosis. (4) Taken patients are separated into 2 groups cally representative according to age,sex,quantity currency of anamnesis, degree of the Kettle index and others.

For this distribution taken to basis that only yardstick index type of the ambience refluxate (A.R). In the 1 group 48 patient's ambiences were acidic (Acidic.  $A=2.6\pm 0.3$ ) and in 28 patients were alkaline ( $Al.A=8.7\pm 0.4$ ) Ph-metric research were carried out by fashion BFRL-S20 ph-meter.

In the process diagnosis from rentgenological, endoscopic, for analyzing function of the external respiration (peakflowmetry and spirometry), common analysis of the blood and deject, necessary times ultrasonography, computer tomography was used. In the preliminary level of the examination through polls-asking determined that is happening with the GERD degree occur of the BOS.

In the next levels of the exploration learned which were injured with GERD and at the same time types of the BOS patient's clinic-diagnosing results (table) Practically healthy 16 people haven't any complaints and objective symptoms of the somatic pathology. Taken facts are recycled used from Student's T-yardstick and diversity of  $P<0.05$  results are acknowledged as reliable.

Results and discussion. The pathology of respiration organs BOS and GERD frequency of occurring together showed the followings. When BA and GERD are diagnosed together in patients Ac.  $A=56,2\%$  and Al.  $A=43,7\%$  ambience refluxates were difned (Difference of statistician symbols  $p<0,05$ ).

Obviously, index of occurring together GERD and BOS according to ambience of refluxate, which patients have acidic ambience (Ac.A) is high. In the GERD occurring degree of the BOS dependence to ambience refluxate in the 2 group patients clinical and endoscopic aspects were learned specially.

The features of clinical and endoscopic symptoms for characterize largely and extensively, primarily, those are essence of the disease that basic symptoms were defined.

Table 1.

The occurring degree the dependence to the ambience of refluxate in the GERD of the BOS symptoms.

Symptoms:	Occurrence degree dependence to the ambience of the refluxate	
	Acidic N=48	Alkaline N=28
1. Heartburn +BOS (Attack asphyxia +chronic cough)	24 (45,8%)	11 (39%)
2. Burp +BOS	17 (35,4%)	6 (21%)
3. Bitter in the <u>mouth</u> +BOS	3 (6,2%)	12(43%)
4. Dysphagia +BOS	18 (37,5%)	11 (39%)
5. Regurgitation + BOS	28 (58,3%)	13 (46,4%)
6. Chronic cough	12 (25%)	3 (11%)
7. Attack asphyxia	18 (37,5%)	6 (21%)

**Note.**  $p < 0,05$  \*  $p < 0,001$  \*\* reliable difference between acidic and alkaline.

Ph-metric results of examination shows that, who are injured with GERD patient's mucus of esophagus and ph-metric index is healthy patients have seriously differences. In the 1 group patients A.R ph-metric indexes are  $2,6 \pm 0,3$ , they differed reliable from index of control group. ( $6,9 \pm 0,8$ ) ( $p < 0,001$ ).

In the 2 group patients this index  $8,7 \pm 0,4$ , they differed from index control ( $p < 0,05$ ) and 1 group ( $p < 0,001$ )

So, developing clinical symptoms and occurring degree of the BOS in the GERD in evaluation dependence calls to the A.R, patients who has Ac. A occurring together BOS is high.

## LITERATURES

1. Баратова Д.С. Особенности клинического течения внепищеводных проявлений гастроэзофагеальной рефлюксной болезни // Молодой ученых. – 2016. – №11. – С. 1139-1141.
2. Белялов Ф.И. Гастроэзофагеальная рефлюксная болезнь: Пособие для врачей. – Иркутск: Иркутск: РИОИГИУВа, 2011. – 23 с.
3. Ивашкин В.Т., Маев И.В., Трухманов А.С. и др. Гастроэзофагеальная рефлюксная болезнь. Клинические рекомендации Российской гастроэнтерологической ассоциации. – М. 2014. – С.23.
4. Ивашкин В.Т., Маев И.В., Трухманов А.С. и др. Клинические рекомендации Российской гастроэнтерологической ассоциации по диагностике и лечению гастроэзофагеальной рефлюксной болезни Российский журнал гастроэнтерологии, гепатологии, колопроктологии (РЖГТК), 2017; 27(4). – С. 75–95.

---

5. Костюкевич О.И., Рылова А.К., Рылова Н.В., Былова Н.А., Колесникова Е.А., Корсунская М.И., Симбирцева А.С. Внепищеводные маски гастроэзофагеальной рефлюксной болезни // РМЖ. Медицинское обозрение. 2018. №7(II). С. 76–81.

6. Yuldasheva D.H. Relationship to the refluxate type of the effectiveness of treatment degree of gastroesophageal reflux disease // European Journal of Research. Austria. Vienna. 2019, № 2. P.110-114.

7. Yuldosheva D.X., Zokirov V.Z., G`ulomova Sh.Q. Non-alcoholic fatty liver: modern view of the problem // Multidisciplinary Peer reviewed journal. India. Vol.6, Issue 12, 2020. – P.286-292.

8. Yuldosheva D.X. Prevalence of non- alcoholic fatty liver disease, clinical and laboratory markers // Eurasian journal of medical and natural sciences. Tashkent. 2022, – P. 94-100.

9. Yuldasheva D. X., Radjabov N. G. Importance of character refluxate in the manifestation of clinical and endoscopic evidence of gastroesophageal reflux disease. // Web of Scientist: International Scientific Research Journal. 2023, Volume-4. №10. P.12-16.